

Fixed-Form Variational Posterior Approximation Through Stochastic Linear Regression

Tim Salimans

Erasmus School of Economics

Erasmus University Rotterdam

salimans@ese.eur.nl

David A. Knowles

University of Cambridge

Abstract

We propose a general algorithm for approximating nonstandard Bayesian posterior distributions. The algorithm minimizes the Kullback-Leibler divergence of an approximating distribution to the intractable posterior distribution. Our method can be used to approximate any posterior distribution, provided that it is given in closed form up to the proportionality constant. The approximation can be any distribution in the exponential family or any mixture of such distributions, which means that it can be made arbitrarily precise. Several examples illustrate the speed and accuracy of our approximation method in practice.

1 Introduction

In Bayesian analysis the posterior distribution is often of non-standard form. To obtain quantities of interest under such a distribution, such as moments or marginal distributions, we typically need to use Monte Carlo methods or approximate the posterior with a more convenient distribution. A popular method of obtaining such an approximation is *structured* or *fixed-form* Variational Bayes, which works by numerically minimizing the Kullback-Leibler divergence of an approximating distribution in the exponential family to the intractable target distribution. For certain problems, algorithms exist that can solve this optimization problem in much less time than it would take to approximate the posterior using Monte Carlo methods (see e.g. Honkela et al., 2010). However, since these methods usually rely on analytic solutions to certain integrals and need conditional conjugacy in the model specification, they are limited in the types of approximations and posteriors they can handle.

We show that solving the optimization problem of fixed-form Variational Bayes is equivalent to performing a linear regression with the sufficient statistics of the approximation as

explanatory variables and the (unnormalized) log posterior density as the dependent variable. Inspired by this result, we present an efficient stochastic approximation algorithm for solving this optimization problem. In contrast to earlier work, our approach does not require any analytic calculation of integrals, which allows us extend the fixed-form Variational Bayes approach to problems where it was previously not applicable. Our method can be used to approximate any posterior distribution, provided that it is given in closed form up to the proportionality constant. The type of approximating distribution can be any distribution in the exponential family or any mixture of such distributions, which means that our approximations can in principle be made arbitrarily precise. While our method somewhat resembles performing stochastic gradient descent on the variational objective function in parameter space, the linear regression view gives insights which allow a more statistically and therefore computationally efficient approach.

Section 2 introduces fixed-form variational posterior approximation, the optimization problem to be solved, and the notation used in the remainder of the paper. In Section 3 we re-interpret this optimization problem as a linear regression problem and we propose a stochastic approximation algorithm to solve it. We suggest several different ways of implementing this algorithm, details of which are given in Sections 4 and 5. In Section 6 we discuss how to assess the quality of the resulting posterior approximations. Here we also present a new method for approximating the marginal likelihood of a model. Our approach allows us to use a larger variety of posterior approximations than was previously possible. Section 7 gives some guidance on how to choose the type of posterior approximation. Here we discuss using mixtures of exponential family distributions and other nonstandard approximations. Section 8 shows some examples of using our method in practice. Here we show that despite its generality, the efficiency of our algorithm is highly competitive with more specialized approaches. Finally, Section 9 concludes.

2 Fixed-form Variational Bayes

Let x be a vector of unknown parameters and/or latent random effects for which we have specified a prior distribution $p(x)$, and let $p(y|x)$ be the likelihood of observing a given set of data y . Upon observing y , we can use Bayes' rule to obtain our updated state of belief, the posterior distribution:

$$p(x|y) = \frac{p(x, y)}{p(y)} = \frac{p(y|x)p(x)}{\int p(y|x)p(x)dx}. \quad (1)$$

An equivalent (Caticha and Giffin, 2006) definition of the posterior distribution is

$$p(x|y) = \arg \min_{q(x)} \mathbb{E}_{q(x)} \log \frac{q(x)}{p(x, y)} = \arg \min_{q(x)} D[q(x)|p(x|y)] - \log p(y), \quad (2)$$

where the optimization is over all proper probability distributions $q(x)$, and where $D[q(x)|p(x|y)]$ denotes the Kullback-Leibler divergence between $q(x)$ and $p(x|y)$. The KL-divergence is always non-negative and has a unique minimizing solution $q(x) = p(x|y)$

almost everywhere, at which point the divergence is zero. Note that the solution of (2) does not depend on the normalizing constant $p(y)$ of the posterior distribution, but that we do obtain it as a by-product of solving $D[q(x)|p(x|y)] = 0$.

The posterior distribution given in (1) is the exact solution of the *variational* optimization problem in (2), but except for certain special cases it is not very useful by itself because it is of non-standard form. This means that we do not have analytical expressions for the posterior moments of x , or for the marginals $p(x_i|y)$ of the multivariate posterior distribution, nor can we determine the normalizing constant $p(y)$. One method of solving this problem is to approximate these quantities using Monte Carlo simulation. A different approach, which we will pursue here, is to restrict the optimization problem in (2) to a reduced set of more convenient distributions Q . If $p(x, y)$ is of conjugate exponential form, choosing Q to be the set of factorized distributions $q(x) = q(x_1)q(x_2) \dots q(x_k)$ often leads to a tractable optimization problem that can be solved efficiently using an algorithm called Variational Bayes Expectation Maximization (VBEM, Beal and Ghahramani, 2002). Such a factorized solution is attractive because it makes the variational optimization problem easy to solve, but it is also very restrictive: it requires a conjugate exponential model and prior specification and it assumes posterior independence between the different blocks of parameters x_i . This means that this factorized approach can be used with few models, and that the solution $q(x)$ may be a poor approximation to the exact posterior (see e.g. Turner et al., 2008).

A different approach to simplifying the variational optimization problem is to restrict the solution set Q to only include distributions of a certain parametric form $q_\eta(x)$, where η denotes the vector of parameters governing the shape of the posterior approximation. This approach is known as *structured* or *fixed-form* Variational Bayes (Honkela et al., 2010; Storkey, 2000; Saul and Jordan, 1996). Usually, the posterior approximation is chosen to be a specific member of the exponential family of distributions:

$$q_\eta(x) = \exp[T(x)\eta - U(\eta)]\nu(x),$$

where $T(x)$ is a $1 \times k$ vector of sufficient statistics, $U(\eta)$ takes care of normalization, and $\nu(x)$ is a base measure. The $k \times 1$ vector η is often called the set of *natural parameters* of the exponential family distribution $q_\eta(x)$. Using this approach, the variational optimization problem in (2) reduces to a parametric optimization problem in η :

$$\hat{\eta} = \arg \min_{\eta} \mathbb{E}_{q_\eta(x)}[\log q_\eta(x) - \log p(x, y)]. \quad (3)$$

If our posterior approximation is of a standard form, the $\mathbb{E}_{q(x)} \log q(x)$ term in (3) can often be evaluated analytically. If we can then also determine $\mathbb{E}_{q(x)} \log p(x, y)$, the optimization problem can be solved using gradient-based optimization or fixed-point algorithms. Posterior approximations of this type are often much more accurate than a factorized approximation, but the requirement that $q_\eta(x)$ is of standard form is still restrictive, as is the requirement of being able to evaluate $\mathbb{E}_{q(x)} \log p(x, y)$. In addition, existing optimization

algorithms for fitting this type of approximation can be much slower than the EM type algorithms used for factorized approximation, reducing somewhat their advantage with respect to Monte Carlo methods. In the next section, we undertake the development of an algorithm that can efficiently solve the variational optimization problem for almost any type of approximating distribution $q_\eta(x)$ and exact posterior $p(x|y)$. The only requirements we impose on $\log p(x, y)$ is that it is given in closed form. The main requirement on $q_\eta(x)$ is that we can sample from it. For simplicity, Sections 3 and 4 will also assume that $q_\eta(x)$ is in the exponential family. Section 7 will then show how we can extend this to include mixtures of exponential family distributions. By using these mixtures and choosing $q_\eta(x)$ to be of a rich enough type, we can in principle make our approximation arbitrarily precise.

3 Variational Bayes as linear regression

For notational convenience we will use an unnormalized version of the approximating distribution:

$$\tilde{q}_{\tilde{\eta}}(x) = \exp[\tilde{T}(x)\tilde{\eta}]\nu(x),$$

where we have removed the normalizer $U(\eta)$, and have added a constant to the vector of sufficient statistics, i.e. $\tilde{T}(x) = (1, T(x))$ and $\tilde{\eta} = (\eta_0, \eta')'$.

The unnormalized version of the KL-divergence is given by

$$D[\tilde{q}_{\tilde{\eta}}(x)|p(x, y)] = \int \tilde{q}_{\tilde{\eta}}(x) \log \frac{\tilde{q}_{\tilde{\eta}}(x)}{p(x, y)} d\nu(x) - \int \tilde{q}_{\tilde{\eta}}(x) d\nu(x) \quad (4)$$

$$= \int \exp[\tilde{T}(x)\tilde{\eta}] [\tilde{T}(x)\tilde{\eta} - \log p(x, y)] d\nu(x) - \int \exp[\tilde{T}(x)\tilde{\eta}] d\nu(x) \quad (5)$$

At the minimum this gives (see Appendix B) $\eta_0 = \mathbb{E}_q \log p(x, y) - \log q(x)$, which is the usual bound on the log evidence. The other parameters, η have the same minimum as in the normalized case.

Taking the gradient of (4) with respect to the natural parameters, $\tilde{\eta}$ we have

$$\nabla_{\tilde{\eta}} D[\tilde{q}_{\tilde{\eta}}(x)|p(x, y)] = \int \tilde{q}_{\tilde{\eta}}(x) [\tilde{T}(x)' \tilde{T}(x) \tilde{\eta} - \tilde{T}(x)' \log p(x, y)] d\nu(x). \quad (6)$$

Setting this expression to zero in order to find the minimum gives

$$\tilde{\eta} = \left[\int \tilde{q}_{\tilde{\eta}}(x) \tilde{T}(x)' \tilde{T}(x) d\nu(x) \right]^{-1} \left[\int \tilde{q}_{\tilde{\eta}}(x) \tilde{T}(x)' \log p(x, y) d\nu(x) \right]. \quad (7)$$

or in its normalized form

$$\tilde{\eta} = [\mathbb{E}_q \tilde{T}(x)' \tilde{T}(x)]^{-1} \mathbb{E}_q \tilde{T}(x)' \log p(x, y) \quad (8)$$

Note that we have implicitly assumed that the Fisher information matrix, $\mathbb{E}_q \tilde{T}(x)' \tilde{T}(x)$ is non-singular, which will be the case for any identifiable approximating exponential family q . Our key insight is to notice the similarity between (8) with the maximum likelihood estimator for linear regression. Recall that in classical linear regression we have that the dependent variable $\{y_n \in \mathbb{R} : n = 1, \dots, N\}$ is distributed as $N(Y|X\beta, \sigma^2 I)$ where X is the $N \times D$ design matrix, β is the $D \times 1$ vector of regression coefficients and σ^2 is the noise variance. The maximum likelihood estimator for β is then

$$\hat{\beta} = (X'X)^{-1}X'Y \quad (9)$$

Alternatively, the estimator in (9) may be seen as the Bayesian posterior mean for β , obtained with the Jeffreys prior on β, σ^2 . Note that this optimal estimator uses the realization of the $X'X$ matrix, rather than its expectation $\mathbb{E}X'X$, even if the latter were available. This observation will be relevant in Section 3.1 where we consider stochastically approximating (8). To see the relation between (8) and (9), associate the design matrix X with the sufficient statistics $\tilde{T}$, the dependent variable Y with the unnormalized log posterior $\log p(x, y)$ and the regression coefficients β with the vector of natural parameters $\tilde{\eta}$. Finally if we consider Monte Carlo estimates of the expectations in (8) then the analogy is very fitting (see (11) below).

3.1 Choosing an estimator

We will consider three different MC estimators for approximating (8). The first separately approximates the two integrals and then calculates the ratio:

$$\hat{\eta}_1 = \left(\frac{1}{S} \sum_r \tilde{T}(x_r)' \tilde{T}(x_r) \right)^{-1} \frac{1}{S} \sum_s \tilde{T}(x_s)' \log p(x_s, y), \quad x_r, x_s \sim_{iid} q(x), \quad (10)$$

with S the number of Monte Carlo samples. The second approximates both integrals using the same samples from q

$$\hat{\eta}_2 = \left(\frac{1}{S} \sum_s \tilde{T}(x_s)' \tilde{T}(x_s) \right)^{-1} \frac{1}{S} \sum_s \tilde{T}(x_s)' \log p(x_s, y), \quad x_s \sim_{iid} q(x) \quad (11)$$

Note that only this estimator is directly analogous to the linear regression estimator. The third estimator is available only when the first expectation is available analytically:

$$\hat{\eta}_a = \left[\mathbb{E}_q \tilde{T}(x)' \tilde{T}(x) \right]^{-1} \frac{1}{S} \sum_s \tilde{T}(x_s)' \log p(x_s, y), \quad x_s \sim_{iid} q(x) \quad (12)$$

We wish to understand the bias/variance tradeoff inherent in each of these estimators. To keep notation manageable consider the case with only $k = 1$ sufficient statistic¹ and let

$$a(x) = \tilde{T}(x)' \tilde{T}(x) = \tilde{T}(x)^2 \quad (13)$$

$$b(x) = \tilde{T}(x) \log p(x, y) \quad (14)$$

¹These results extend in a straightforward if laborious manner to the case where $k > 1$

We can now write the three estimators of η more concisely as

$$\hat{\eta}_1 = \frac{\frac{1}{S} \sum_r b(x_r)}{\frac{1}{S} \sum_s a(x_s)}, \quad x_r, x_s \sim_{iid} q(x) \quad (15)$$

$$\hat{\eta}_2 = \frac{\frac{1}{S} \sum_s b(x_s)}{\frac{1}{S} \sum_s a(x_s)}, \quad x_s \sim_{iid} q(x) \quad (16)$$

$$\hat{\eta}_a = \frac{\frac{1}{S} \sum_s b(x_s)}{\mathbb{E}a}, \quad x_s \sim_{iid} q(x) \quad (17)$$

Using a simple Taylor series argument it is straightforward (see Appendix A) to approximate the bias of these estimators:

$$\text{bias}(\hat{\eta}_1) \approx \frac{\text{Var}(a)\mathbb{E}[b]}{S\mathbb{E}[a]^3} \quad (18)$$

$$\text{bias}(\hat{\eta}_2) \approx \frac{\text{Var}(a)\mathbb{E}[b]}{S\mathbb{E}[a]^3} - \frac{\text{Cov}(a,b)}{S\mathbb{E}[a]^2} \quad (19)$$

Note that the first term is shared, but the first estimator does not have the covariance term as a result of the independent sampling in approximating the numerator and denominator. In contrast $\hat{\eta}_a$ is unbiased. Now consider the variances

$$\text{Var}(\hat{\eta}_1) \approx \frac{1}{S} \left(\frac{(\mathbb{E}b)^2 \text{Var}(a)}{(\mathbb{E}a)^4} + \frac{\text{Var}(b)}{(\mathbb{E}a)^2} \right) \quad (20)$$

$$\text{Var}(\hat{\eta}_2) \approx \frac{1}{S} \left(\frac{(\mathbb{E}b)^2 \text{Var}(a)}{(\mathbb{E}a)^4} - 2 \frac{\mathbb{E}b \text{Cov}(a,b)}{(\mathbb{E}a)^3} + \frac{\text{Var}(b)}{(\mathbb{E}a)^2} \right) \quad (21)$$

$$\text{Var}(\hat{\eta}_a) = \frac{\text{Var}(b)}{S(\mathbb{E}a)^2} \quad (22)$$

All three estimators have the same final term (the variance of the “analytic” estimator). Again the second estimator has an additional term resulting from the covariance between a and b which we find is typically beneficial in that it results in the variance of $\hat{\eta}$ being significantly smaller. It is worth recalling that the mean squared error (MSE) of an estimator is given by

$$\mathbb{E}[(\eta - \hat{\eta})^2] = \text{Var}(\hat{\eta}) + \text{bias}(\hat{\eta})^2 \quad (23)$$

Since both the variance and bias are $O(1/S)$, the variance contribution to the MSE is $O(1/S)$ whereas the bias contribution is $O(1/S^2)$, so the variance is actually a greater problem than the bias. From these expressions it is still not immediately obvious which estimator we should use. However, consider the case when the target distribution p is in the same exponential family as q , i.e. when $\log p(x, y) = \tilde{T}(x)\lambda$. It is then straightforward to show that

$$\text{bias}(\hat{\eta}_1) \approx \frac{\lambda \text{Var}(\tilde{T}^2)}{S\mathbb{E}[\tilde{T}^2]^2}, \quad \text{Var}(\hat{\eta}_1) \approx 2 \frac{\lambda^2 \text{Var}(\tilde{T}^2)}{S\mathbb{E}[\tilde{T}^2]^2} \quad (24)$$

$$\text{bias}(\hat{\eta}_2) \approx 0, \quad \text{Var}(\hat{\eta}_2) \approx 0 \quad (25)$$

$$\text{bias}(\hat{\eta}_a) = 0, \quad \text{Var}(\hat{\eta}_a) = \frac{\lambda^2 \text{Var}(\tilde{T}^2)}{S\mathbb{E}[\tilde{T}^2]^2} \quad (26)$$

We see that in this case for $\hat{\eta}_2$ the positive and negative contributions to both the bias and variance cancel. While this result will not hold exactly for cases of interest, it suggests that for exponential families which are capable of approximating p reasonably well, $\hat{\eta}_2$ should perform significantly better than $\hat{\eta}_1$ or even $\hat{\eta}_a$. In this setting it is actually possible to see that $\hat{\eta}_2$ will in fact give the exact solution in $k + 1$ samples (with k the number of sufficient statistics), while the other estimators have non-vanishing variance for a finite number of samples. This means that the approximate equality in (25) can be replaced by exact equality. Using $k + 1$ samples $x_i, i = 1, \dots, k + 1$, assumed to be unique (which holds almost surely for continuous distributions q), we have

$$\hat{\eta}_2 = \left(\sum_{i=1}^{k+1} \tilde{T}(x_i)' \tilde{T}(x_i) \right)^{-1} \sum_{i=1}^{k+1} \tilde{T}(x_i)' \tilde{T}(x_i) \lambda = \lambda \quad (27)$$

That is, the algorithm has recovered $p(x, y)$ exactly with probability one. If we assume we know how to normalize q , this means we also have $p(x|y)$ exactly in this case. Note that we recover the exact answer here because the $p(x, y)$ function evaluations are in themselves *noise free*, so the regression analogy really corresponds to a noise free regression.

It is instructive to consider a toy example: fitting an exponential distribution $p(x) = \lambda e^{-\lambda x}$, about the simplest possible demonstration of the exact fitting phenomenon shown in (27). We assume that we are unaware that p happens to be normalized. Our variational approximation has $\tilde{T} = [1, x]'$ and rate η , i.e. $q(x) = \eta e^{-\eta x}$. Note that is an example where it is straightforward to calculate

$$\mathbb{E}_q \tilde{T}(x)' \tilde{T}(x) = \begin{bmatrix} 1 & -\eta^{-1} \\ -\eta^{-1} & \eta^{-2} \end{bmatrix}$$

We test the three estimators in (10), (11) and (12) when the true exponential rate is $\lambda = 1.5$, and sampling from the optimal q distribution with $\eta = 1.5$. The results confirm that $\hat{\eta}_2$ finds the exact rate using just $S = 2$ MC samples, as predicted by (27). We would expect $\hat{\eta}_a$ to be unbiased, and this is borne out by the results shown in Figure 1. The estimator $\hat{\eta}_1$ has both poor bias and such large variance that it often gives an invalid negative rate if fewer than 10 MC samples are used. While this is clearly a very simple example it hopefully emphasizes the potential benefit to be gained from using estimators related to $\hat{\eta}_2$.




Figure 1: Comparison of three estimators for fitting a variational posterior q to a simple exponential distribution p . 50 repeats were used to estimate the mean and variance of the estimator: the thick line shows the mean and the thin lines show $\pm$ one standard deviation. The x -axis indicator the number of MC samples, S , used. As expected in this case $\hat{\eta}_2$ gives the correct solution of 1.5 using $S \geq 2$ samples.

3.2 A stochastic approximation algorithm

We have seen that the linear regression view of variational inference inspires a statistically efficient estimator for the natural parameters of the variational approximation in the form of (11). The question remains how we should turn this estimator into an automated algorithm. Note that in the general case where q and p are of different forms, the linear regression in equation (8) only gives the optimal $\hat{\eta}$ parameters when we sample from the optimal approximation $q_{\hat{\eta}}$. Since we do not know this optimal approximation before having done the optimization, this creates a chicken-and-egg problem. The obvious solution would be to use an arbitrary initial distribution q , and use the optimality condition in (8) as a fixed-point equation to be iterated until converge, which would give updates of the form

$$\eta_{t+1} = [\mathbb{E}_{q(\eta_t)} \tilde{T}(x)' \tilde{T}(x)]^{-1} \mathbb{E}_{q(\eta_t)} \tilde{T}(x)' \log p(x, y). \quad (28)$$

However, such an approach is not guaranteed to converge, and in practice we find it often does not. Fortunately, the difference $\eta_{t+1} - \eta_t$ generated by this approach can be interpreted as the negative approximate natural gradient of the Kullback-Leibler divergence (4) (see Appendix D). This means that we can obtain a convergent algorithm by taking small enough steps in the direction indicated by (28).

Based on this insight, we propose solving the variational optimization problem in (8) using the following stochastic optimization algorithm. The basic idea is to sample one MC

point at a time, updating the current estimate of q as our estimate of $\hat{\eta}$ improves. Thus samples drawn later in the process will be closer to being samples from the optimal q . This approach allows us to efficiently approximate the statistics used in the regression of (8), while adapting η slowly enough to ensure convergence of the algorithm.

Algorithm 1 Stochastic Optimization for Fixed-Form Variational Bayes

Require: An unnormalized posterior distribution $p(x, y)$

Require: A type of approximating posterior $q_\eta(x)$

Require: The total number of iterations N

Initialize η to a first guess, for example by matching the prior $p(x)$

Initialize $C = \mathbb{E}_{q_\eta} \tilde{T}(x)' \tilde{T}(x)$, or a diagonal approximation of this matrix

Initialize $g = C\eta$

Initialize $\bar{C} = \mathbf{0}$

Initialize $\bar{g} = \mathbf{0}$

Step-size $w = 1/\sqrt{N}$

for $t = 1 : N$ **do**

Set $\eta = C^{-1}g$

Generate $\hat{g}_t =$ unbiased estimate of $\mathbb{E}_{q_\eta} \tilde{T}(x)' \log p(x, y)$

Generate $\hat{C}_t =$ unbiased estimate of $\mathbb{E}_{q_\eta} \tilde{T}(x)' \tilde{T}(x)$

Set $g = (1 - w)g + w\hat{g}_t$

Set $C = (1 - w)C + w\hat{C}_t$

if $t > N/2$ **then**

Set $\bar{g} = \bar{g} + \hat{g}_t$

Set $\bar{C} = \bar{C} + \hat{C}_t$

end if

end for

return $\hat{\eta} = \bar{C}^{-1}\bar{g}$

This algorithm is inspired by a long line of research on stochastic approximation, starting with the seminal work of Robbins and Monro (1951). However, contrary to most applications in the literature, we use a fixed step size $w = 1/\sqrt{N}$ rather than a declining one in updating our statistics. The analyses of Robbins and Monro (1951) and Amari (1997) show that a sequence of learning rates $w_t = ct^{-1}$ is asymptotically efficient in stochastic gradient descent as the number of iterations N goes to infinity, but this conclusion rests on very strong assumptions on the functional form of the objective function (e.g. strong convexity) that are not satisfied for the problems we are interested in. Moreover, with a finite number of iterations N , the effectiveness of a sequence of learning rates that decays this fast is highly dependent on the proportionality constant c . If we choose c either too low or too high, it may take an extremely long time to reach the efficient asymptotic regime of this learning rate sequence. Nemirovski et al. (2009) show that a more robust approach is to use a constant learning rate $w = 1/\sqrt{N}$ and that this is optimal for finite N without putting stringent requirements on the objective function. In order to reduce the

variance of the last iterate with this non-vanishing learning rate, they propose to use an average of the last L iterates as the final output of the optimization algorithm. The value of L should grow with the total number of iterations, and is usually chosen to be equal to $N/2$. Remarkably, they show that such an averaging procedure can match the asymptotic efficiency of the optimal learning sequence $w_t = ct^{-1}$. For our particular optimization problem we have observed excellent results using constant learning rate $w = 1/\sqrt{N}$, and averaging starting half-way into the optimization. Note that we perform this averaging on the statistics g and C , rather than on the parameters $\eta = C^{-1}g$, which is statistically more efficient for our application. Using this set-up, g and C are actually weighted MC estimates where the weight of the j -th MC sample during the t -th iteration ($j \leq t$) is given by $w(1-w)^{t-j}$. Since $w \in (0, 1)$, this means that the weight of earlier MC samples declines as the algorithm advances, which is desirable since we expect q to be closer to optimal later in the algorithm's progression.

If the initial guess for η is very far from the optimal value, or if the number of steps N is very small, it can sometimes occur that the algorithm proposes a new value for η that does not define a proper distribution. We can guard against this by adapting the regression step $\eta = C^{-1}g$ to have additional shrinkage towards the current value of η by using

$$\eta \leftarrow (C + \Lambda)^{-1}(g + \Lambda\eta),$$

with Λ a positive diagonal matrix of appropriate scale. The scale of Λ can be made to depend on the sampled $\hat{g}_t, \hat{C}_t$ to prevent very large moves in η : as long as the sampled statistics are still averaged in the usual way this does not bias the algorithm.

Alternatively we could just reduce the step-size or increase the number of steps: One can show that the algorithm above becomes a pre-conditioned gradient descent algorithm as the number of steps goes to infinity (see Appendix D), which means that the algorithm is guaranteed not to diverge if the step size is small enough. In addition, note that the exact convergence result of equation (27) suggests that divergence is very unlikely if $q_\eta(x)$ and $p(x, y)$ are close in functional form: choosing a good type of approximation will thus also help to ensure fast convergence. Finally, picking a good first guess for η also helps the algorithm to converge more quickly. For very difficult cases it might therefore be worthwhile to base this guess on a first rough approximation of the posterior, for example by choosing η to match the curvature of $\log p(x, y)$ at its mode. For all our applications we found that a simple first guess for η and a large enough number of iterations was sufficient to guarantee a stable algorithm.

Finally, it is worth mentioning the calculation of $C^{-1}g$. Note that by using conditional independencies in the exact posterior (Section 4.2) and the structure of the approximation (Section 5), we can often avoid computing and storing the full C matrix, which can be quite high dimensional. In those rare cases where we do use the full C matrix, computing C^{-1} explicitly (which is $\mathcal{O}(K^3)$) is not recommended, but if desired then C^{-1} should be updated each iteration using rank-one updates (i.e. using the matrix inversion lemma) which cost

$\mathcal{O}(K^2)$. Similar low cost updates could be used to maintain Cholesky decompositions since C is symmetric, which is a numerically stable and efficient option. In very high dimensions one could use conjugate gradients to solve $C^{-1}g$ approximately.

4 Simulating the regression statistics

In order to implement our algorithm we need to be able to construct unbiased estimates of $\mathbb{E}_{q_\eta} \tilde{T}(x)' \log p(x, y)$ and $\mathbb{E}_{q_\eta} \tilde{T}(x)' \tilde{T}(x)$. It turns out that this is easy to do as long as $q_\eta(x)$ is easy to sample from. In fact, there are multiple ways of constructing stochastic estimates $\hat{g}_t$ and $\hat{C}_t$ with these expectations. We will derive several here.

4.1 Basic stochastic approximation

Following the results of Section 3.1 our default stochastic approximate of $\mathbb{E}_{q_\eta} \tilde{T}(x)' \log p(x, y)$ and $\mathbb{E}_{q_\eta} \tilde{T}(x)' \tilde{T}(x)$ is to simply draw a sample x^* from $q_\eta(x)$ and to use this sample to calculate

$$\hat{g}_t = \tilde{T}(x^*)' \log p(x^*, y) \quad (29)$$

$$\hat{C}_t = \tilde{T}(x^*)' \tilde{T}(x^*) \quad (30)$$

This works remarkably well because, as Section 3.1 explains, using the same random draw x^* to form both estimates, part of the random variation in $\eta = C^{-1}g$ cancels out.

4.2 Making use of conditional independencies

For most statistical problems, the log posterior can be decomposed into a number of additive factors, i.e. $\log p(x, y) = \sum_{j=1}^N \log p_j(x, y)$. The optimality condition in equation (8) can then also be written as a sum:

$$\tilde{\eta} = \sum_{j=1}^N [\mathbb{E}_q \tilde{T}(x)' \tilde{T}(x)]^{-1} \mathbb{E}_q \tilde{T}(x)' \log p_j(x, y)$$

This means that rather than performing one single linear regression we can equivalently perform N separate regressions.

$$\hat{\eta} = \sum_{j=1}^N \hat{\eta}^j \quad (31)$$

$$\hat{\eta}^j = [\mathbb{E}_q \tilde{T}(x)' \tilde{T}(x)]^{-1} \mathbb{E}_q \tilde{T}(x)' \log p_j(x, y) \quad (32)$$

The practical benefit of this is that these separate regressions are often of much lower dimension: We know that element i of $\hat{\eta}_j$ will only be non-zero if the i -th sufficient statistic $\tilde{T}_i(x)$ has non-zero partial correlation to $\log p_j(x, y)$. Since the separate factors $\log p_j(x, y)$

often involve only a subset of the variables in x , this means that we can omit many of the sufficient statistics in performing each regression. That is, we have

$$\hat{\eta}_R^j = [\mathbb{E}_q \tilde{T}_R(x)' \tilde{T}_R(x)]^{-1} \mathbb{E}_q \tilde{T}_R(x)' \log p_j(x, y)$$

with $\tilde{T}_R(x)$ the relevant subset of $\tilde{T}(x)$, and $\hat{\eta}_R^j$ the corresponding subset in $\hat{\eta}^j$. The remaining elements in $\hat{\eta}^j$ will be zero. By performing these lower dimensional regressions we can reduce the variance of the stochastic approximation algorithm, as well as reduce the overhead needed to store and invert $\mathbb{E}_q \tilde{T}(x)' \tilde{T}(x)$.

4.3 Using the gradient of the log posterior

It is straightforward to split the optimality condition in (8) into a condition for the normalizer η_0 and for the original vector of natural parameters η . Following Appendix B, the optimality condition for the original parameters of the normalized q -distribution is given by

$$\hat{\eta} = \text{Cov}_q[T(x), T(x)]^{-1} \text{Cov}_q[T(x), \log p(x, y)]. \quad (33)$$

Furthermore, using the properties of the exponential family of distributions, we know that

$$\text{Cov}_q[T(x), T(x)] = \nabla_\eta \mathbb{E}_{q_\eta} T(x) \quad (34)$$

and

$$\text{Cov}_q[T(x), \log p(x, y)] = \nabla_\eta \mathbb{E}_{q_\eta} \log p(x, y) \quad (35)$$

Both expectations can be approximated without bias using Monte Carlo. By differentiating these Monte Carlo approximations we can then obtain unbiased estimates of their derivatives. This is easy to do as long as the pseudo-random draw x^* from $q_\eta(x)$ is a differentiable function of the parameters η , given our random number seed z^* .

$$x^* = f(\eta, z^*), \text{ with } z^* \text{ such that } x^* \sim q_\eta(x) \quad (36)$$

$$\hat{g} = \nabla_\eta \log p(f(\eta, z^*), y) \quad (37)$$

$$\hat{C} = \nabla_\eta T(f(\eta, z^*)) \quad (38)$$

By using the same random number seed z^* in both Monte Carlo approximations we once again get the beneficial variance reduction effect described in Section 3.1. Empirically, we find that using gradients often leads to a more efficient stochastic optimization algorithm. For some applications the posterior distribution will not be differentiable in some of the elements of x , for example when x is discrete. In that case the stochastic approximations presented here may be combined with those of Section 3.1.

Furthermore, we can decompose $\nabla_\eta \log p(f(\eta, z^*), y) = \nabla_\eta f(\eta, z^*) \nabla_x \log p(x, y)$ evaluated at $x = f(\eta, z^*)$, and equivalently $\nabla_\eta T(f(\eta, z^*)) = \nabla_\eta f(\eta, z^*) \nabla_x T(x)$. Note that for many

samplers $\nabla_\eta f(\eta, z^*)$ is not defined, e.g. rejection samplers. In that case we can replace this quantity by the equivalent expression for a sample from an inverse-transform sampler:

$$\nabla_\eta f(\eta, z^*) = -\frac{\nabla_\eta \Phi_\eta(x^*)}{\phi_\eta(x^*)}$$

with $\Phi_\eta(x^*)$ the CDF and $\phi_\eta(x^*)$ the pdf of the sampler.

4.4 Using the Hessian of the log posterior

When we have both first and second order gradient information for $\log p(x, y)$ and if we choose our approximation to be multivariate Gaussian, i.e. $q_\eta(x) = N(m(\eta), V(\eta))$, we have a third option for approximating the statistics used in the regression. For Gaussian $q(x)$ and twice differentiable $\log p(x, y)$, Minka (2001b) and Opper and Archambeau (2009) show that

$$\nabla_m \mathbb{E}_q \log p(x, y) = \mathbb{E}_q \nabla_x \log p(x, y) \quad (39)$$

and

$$\nabla_V \mathbb{E}_q \log p(x, y) = \frac{1}{2} \mathbb{E}_q \nabla_x \nabla_x \log p(x, y) \quad (40)$$

where $\nabla_x \nabla_x \log p(x, y)$ denotes the Hessian matrix of $\log p(x, y)$ in x .

For the multivariate Gaussian distribution we know that the natural parameters are given as $\eta_1 = V^{-1}m$ and $\eta_2 = V^{-1}$. Using this relationship, we can derive Monte Carlo estimators $\hat{g}$ and $\hat{C}$ using the identities (34,35). We find that these stochastic approximations are often even more efficient than the ones in Section 4.3, provided that the Hessian matrix of $\log p(x, y)$ can be calculated cheaply.

4.5 Using analytic expectations where possible

In many cases it is possible to calculate the contributions of some of the factors $\log p_i(x)$ to the stochastic approximations C and g analytically, while for others it is not. For example, this occurs when part of $\log p(x)$ (most often the prior) is conjugate to the posterior approximation $q_\eta(x)$. Even for some non-conjugate factors it might be possible to calculate certain expectations analytically. Using these exact expectations rather than their stochastic estimates can help reduce the variance of the approximations as well as reduce the time required to compute them, both of which increase the efficiency of the optimization procedure.

4.6 Subsampling the data: double stochastic approximation

The stochastic approximations derived above are all linear functions of $\log p(x)$ and its first and second derivatives. This means that these estimates are still unbiased even if we take

$\log p(x)$ to be a noisy unbiased estimate of the true log posterior, rather than the exact log posterior. For most statistical applications $\log p(x)$ itself is a separable additive function of a number of independent factors, i.e. $\log p(x) = \sum_{i=1}^N \log p_i(x)$. These $\log p_i(x)$ terms can be the likelihood contributions of individual observed data points, but they can also arise through conditional independencies between the x variables in the posterior. Using this fact we can construct an unbiased stochastic approximation of $\log p(x)$ as

$$\log \tilde{p}(x) = \frac{N}{K} \sum_{j=1}^K \log p_j(x) \quad (41)$$

where the K factors $\log p_j(x)$ are randomly selected from the total N factors. This approach was previously proposed for online learning of topic models by Hoffman et al. (2010). Since $\log \tilde{p}(x)$ has $\log p(x)$ as its expectation, performing stochastic approximation based on $\tilde{p}(x)$ converges to the same solution as when using $p(x)$, provided we resample the factors in $\log \tilde{p}(x)$ at every iteration. By subsampling the $K \ll N$ factors in the model the individual steps of the optimization procedure become more noisy, but since we can calculate $\tilde{p}(x)$ faster than we can $p(x)$, we can perform a larger number of steps in the same amount of time. If the number of factors in the posterior is especially large, this tradeoff often favors using subsampling. This principle has been used in many successful applications of stochastic gradient descent, see e.g. Bottou (2010).

5 Linear transformations of the regression problem

It is well known that classical linear least squares regression is invariant to invertible linear transformations of the explanatory variables. More precisely, when we have an $N \times D$ matrix of explanatory variables X , and an $N \times 1$ vector of dependent variables Y , we may equivalently perform the linear regression using a transformed set of explanatory variables $\tilde{X} = XK'$, with K any invertible matrix of size $D \times D$. The least squares estimator $\tilde{\beta} = (\tilde{X}'\tilde{X})^{-1}\tilde{X}'Y$ of the transformed problem can then be used to give the least squares estimator of the original problem as $\hat{\beta} = K'\tilde{\beta}$:

$$\hat{\beta} = K'(KX'XK')^{-1}KX'Y = (KX'X)^{-1}KX'Y = (X'X)^{-1}X'Y.$$

Using the same principle, we can rewrite the optimality condition of equation (8) as

$$\tilde{\eta} = [\mathbb{E}_{q_\eta} K(\eta) \tilde{T}(x)' \tilde{T}(x)]^{-1} \mathbb{E}_{q_\eta} K(\eta) \tilde{T}(x)' \log p(x, y) \quad (42)$$

for any invertible matrix K , which may depend on the variational parameters η . Instead of solving our original least squares regression problem, we may thus equivalently solve a transformed version of it as defined by any invertible linear transformation $K(\eta)$. When we perform the linear regression of equation (42) for a fixed set of parameters η , the result will obviously be identical to that of the original regression with $K(\eta) = \mathbf{I}$, as long as we use the same random numbers for both regressions. However, when the ‘data points’ in

our regression are generated using different values of η , as is the case with the proposed stochastic approximation algorithm, the two regressions will not necessarily give the same solution. If the true posterior $p(x|y)$ is of the same functional form as the approximation q_η , the exact convergence result of Section 3 holds for any invertible $K(\eta)$, so it is not immediately obvious which $K(\eta)$ is best for general applications.

We hypothesize that certain choices of $K(\eta)$ may lead to statistically more efficient stochastic approximation algorithms for certain specific problems, but we will not pursue this idea here. What we will discuss is the observation that the stochastic approximation algorithm may be easier to implement for some choices of $K(\eta)$ than for others, and that the computational costs are not identical for all $K(\eta)$. In particular, it is worth noting that the transformation $K(\eta)$ allows us to use different parameterizations of the variational approximation. Let q_ϕ be such a reparameterization of the approximation, let the new parameter vector $\phi(\eta)$ be an invertible and differentiable transformation of the original parameters η , and set $K(\eta)$ equal to the inverse Jacobian of this transformation, i.e. $K(\eta) = [\nabla_\eta \phi(\eta)]^{-1}$. Using the properties of the exponential family of distributions, we can then show that

$$K(\eta) \text{Cov}_{q_\phi}[T(x), h(x)] = \nabla_\phi \mathbb{E}_{q_\phi} h(x) \quad (43)$$

for any differentiable function $h(x)$. Using this result, the stochastic approximations of Section 4.3 for the transformed regression problem are found to be

$$x^* = f(\phi, z^*), \text{ with } z^* \text{ such that } x^* \sim q_\phi(x) \quad (44)$$

$$\hat{g} = \nabla_\phi \log p(f(\phi, z^*), y) \quad (45)$$

$$\hat{C} = \nabla_\phi T(f(\phi, z^*)). \quad (46)$$

These new expressions for $\hat{g}$ and $\hat{C}$ may be easier to calculate than the original ones (36), and the resulting $\hat{C}$ may have a structure making it easier to invert in some cases. A particularly striking example of this occurs when we use a Gaussian approximation in combination with the stochastic approximations of Section 4.4, using the gradient and Hessian of $\log p(x, y)$. In this case we may work in the usual natural parameterization, but doing so gives a dense matrix $\hat{C}$ with dimensions proportional to p^2 , where p is the dimension of x . For large p , such a stochastic approximation is expensive to store and invert. However, using the stochastic approximations above, we may also parameterize our approximation in terms of the mean m and variance V , and work with these parameters directly. Doing so leads to the following sparse regression algorithm, as derived in Appendix C.

Algorithm 2 Stochastic Approximation for Gaussian Variational Approximation

Require: An unnormalized, twice differentiable posterior distribution $p(x, y)$

Require: The total number of iterations N

Initialize the mean and variance of the approximation (m, V) to a first guess, for example by matching the prior $p(x)$

Initialize $z = m$, $P = V^{-1}$ and $a = 0$

Initialize $\bar{z} = 0$, $\bar{P} = \mathbf{0}$ and $\bar{a} = \mathbf{0}$

Step-size $w = 1/\sqrt{N}$

for $t = 1 : N$ **do**

 Set $V = P^{-1}$ and $m = Va + z$

 Generate a draw x^* from $N(m, V)$

 Calculate the gradient g_t and Hessian H_t of $\log p(x, y)$ at x^*

 Set $a = (1 - w)a + wg_t$

 Set $P = (1 - w)P - wH_t$

 Set $z = (1 - w)z - wx^*$

if $t > N/2$ **then**

 Set $\bar{a} = \bar{a} + g_t$

 Set $\bar{P} = \bar{P} - H_t$

 Set $\bar{z} = \bar{z} + x^*$

end if

end for

Set $V = \bar{P}^{-1}$ and $m = V\bar{a} + \bar{z}$

return m, V

Instead of storing and inverting the full C matrix, this algorithm uses the sparsity induced by the transformation $K(\eta)$ to work with the precision matrix P instead. The dimensions of this square matrix are equal to p , rather than its square, providing great savings. Moreover, while the C matrix in the original parameterization is always dense, P will have the same sparsity pattern as the Hessian of $\log p(x, y)$, which may reduce the costs of storing and inverting it even further for many applications. An example of this algorithm can be found in Section 8.1.

6 Marginal likelihood and approximation quality

To fit our posterior approximation we minimize the Kullback-Leibler divergence between $q_\eta(x)$ and $p(x|y)$, given by

$$D(q_\eta|p) = \mathbb{E}_{q_\eta} \log \frac{q_\eta(x)}{p(x|y)} = \mathbb{E}_{q_\eta} \left[\log \frac{q_\eta(x)}{p(x, y)} \right] + \log p(y),$$

which shows that we need to know the marginal likelihood $p(y)$ (the normalizing constant of the posterior) in order to evaluate this Kullback-Leibler divergence. As discussed before,

we do not need to know this constant in order to minimize $D(q_\eta|p)$ as $p(y)$ does not depend on η , but we do need know it if we want to determine the quality of the approximation, as measured by the final KL-divergence. In addition, the constant $p(y)$ is also essential for performing Bayesian model comparison or model averaging.

When our algorithm has converged, we have the following identity

$$\log p(x, y) = \hat{\eta}_0 + \log q_\eta(x) + r(x),$$

where $r(x)$ is the ‘residual’ or ‘error term’ in the linear regression of $\log p(x, y)$ on the sufficient statistics of $q_\eta(x)$. The intercept of the regression is

$$\hat{\eta}_0 = \mathbb{E}_{q_\eta} [\log p(x, y) - \log q_\eta(x)],$$

the usual VB lower bound on the marginal likelihood. Exponentiating this term yields

$$p(x, y) = \exp(\hat{\eta}_0) q_\eta(x) \exp(r(x)),$$

which we need to integrate with respect to x in order to find the marginal likelihood $p(y)$. Doing so gives

$$p(y) = \exp(\hat{\eta}_0) \mathbb{E}_{q_\eta} \exp(r(x)). \quad (47)$$

At convergence we have that $\mathbb{E}_{q_\eta} r(x) = 0$. Jensen’s inequality then tells us that

$$\mathbb{E}_{q_\eta} \exp(r(x)) \geq 1,$$

which shows that $\hat{\eta}_0$ is indeed a lower bound on the log marginal likelihood as we claimed earlier. If our approximation is perfect, the KL-divergence is zero and $r(x)$ is zero almost everywhere. In that case the residual term vanishes and the lower bound will be tight, otherwise it will underestimate the true marginal likelihood. The lower bound $\hat{\eta}_0$ is often used in model comparison, which works well if the KL-divergence between the approximate and true posterior distribution is of approximately the same size for all models that are being compared. However, if we compare two very different models this will often not be the case, and the model comparison will be biased as a result. In addition, as opposed to the exact marginal likelihood, the lower bound gives us no information on the quality of our posterior approximation. It would therefore be useful to obtain a better estimate of the marginal likelihood. One approach to doing this would be to evaluate the expectation in (47) using Monte Carlo sampling. Some analysis shows that this corresponds to approximating $p(y)$ using importance sampling, with $q_\eta(x)$ as the candidate distribution. It is well known that this estimator of the marginal likelihood may have infinite variance, unless $r(x)$ is bounded from above. In general, we cannot guarantee the boundedness of $r(x)$ for our approach, so we will instead approximate the expectation in (47) using something that is easier to calculate. At convergence, we know that the mean of $r(x)$ is zero when sampling from $q_\eta(x)$. The variance of $r(x)$ can easily be estimated using the mean squared error

of the regressions we perform during the optimization, with relatively low variance. We denote our estimate of this variance by s^2 . The assumption we will then make in order to approximate $\log p(y)$ is that $r(x)$ is approximately distributed as a normal random variable with these two moments. This leads to the following simple estimate of the log marginal likelihood

$$\log p(y) \approx \hat{\eta}_0 + \frac{1}{2}s^2.$$

That is, our estimate of the marginal likelihood is equal to its lower bound plus a correction term that captures the error in our posterior approximation $q_\eta(x)$. Similarly, we can approximate the KL-divergence of our posterior approximation as

$$D(q_\eta|p) \approx \frac{1}{2}s^2.$$

The KL-divergence is approximately equal to half the mean squared error in the regression of $\log p(x, y)$ on the sufficient statistics of the approximation. This relationship should not come as a surprise: this mean squared error is exactly what we minimize when we do a linear regression. Our experiments indicate that this approximation of the KL-divergence can be quite accurate (see Section 8.3), especially when the approximation $q_\eta(x)$ is reasonably good.

Note that the scale of the KL-divergence is highly dependent on the amount of curvature in $\log p(x|y)$ and is therefore not easily comparable across different problems. If we scale the approximate KL-divergence to account for this curvature, this naturally leads to the R-squared measure of fit for regression modeling:

$$R^2 = 1 - \frac{s^2}{\text{Var}_q[\log p(x, y)]}$$

The R-squared measure corrects for the amount of curvature in the posterior distribution and is therefore comparable across different models and data sets. In addition it is a well-known measure and easily interpretable. We therefore propose to use the R-squared as the measure of approximation quality for our variational posterior approximations.

7 Using mixtures of exponential family distributions

So far, we have assumed that the approximating distribution $q_\eta(x)$ is a member of the exponential family. Here we will relax that assumption. If we choose a non-standard approximation, this most likely means that certain moments or marginals of $q_\eta(x)$ are no longer available analytically, which should be taken into account when choosing the type of approximation. However, if we can at least sample directly from $q_\eta(x)$, it is often still much cheaper to approximate these moments using Monte Carlo than it would be to approximate the corresponding moments of $p(x|y)$ using MCMC or other indirect sampling methods. We have identified two general strategies for constructing useful non-standard posterior approximations which are discussed in the following two sections.

7.1 Hierarchical approximations

If we split our vector of unknown parameters x into p non-overlapping blocks, our approximating posterior may be decomposed as

$$q(x) = q(x_1)q(x_2|x_1)q(x_3|x_2, x_1) \dots q(x_p|x_{p-1}, \dots, x_1).$$

If we then choose every conditional posterior $q(x_i|x_{i-1}, x_{i-2}, \dots, x_1)$ to be of a standard form, we can easily sample from the joint $q(x)$, while still having much more freedom in capturing the dependence between the different blocks of x . In practice, such a conditionally standard approximation can be achieved by specifying the sufficient statistics of each standard block $q(x_i|x_{i-1}, x_{i-2}, \dots, x_1)$ to be a function of the preceding elements $x_{i-1}, x_{i-2}, \dots, x_1$. This leads to a natural type of approximation for hierarchical Bayesian models, where the hierarchical structure of the prior often suggests a good hierarchical structure for the posterior approximation.

If every conditional $q(x_i|x_{i-1}, x_{i-2}, \dots, x_1)$ is in the exponential family, the joint may not be if the normalizing constant of $q(x_i|x_{i-1}, x_{i-2}, \dots, x_1)$ is a non-separable function of $x_{i-1}, x_{i-2}, \dots, x_1$ and the variational parameters. However, because the conditionals are still in the exponential family, our optimality condition still holds separately for the variational parameters of each conditional with only slight modification. In that case we therefore propose applying the optimization procedures separately to each of these conditionals. Without loss of generalization, consider the case where our posterior approximation consists of two factors: $q(x) = q_{\eta_1}(x_1)q_{\eta_2}(x_2|x_1)$. In its normalized form (see Section 4.3), the optimality condition for the first factor is then given as

$$\eta_1 = \text{Var}_q[T(x_1)]^{-1} \text{Cov}_q[T(x_1), \log p(x, y) - \log q_{\eta_2}(x_2|x_1)],$$

where $T(x_1)$ denotes the sufficient statistics of $q_{\eta_1}(x_1)$. The optimality condition for the second block is

$$\eta_2 = [\mathbb{E}_{q(x_1)} \text{Var}_{q(x_2|x_1)}(T(x_2))]^{-1} [\mathbb{E}_{q(x_1)} \text{Cov}_{q(x_2|x_1)}(T(x_2), \log p(x, y) - \log q_{\eta_1}(x_1))],$$

where $T(x_2)$ denotes the sufficient statistics of $q_{\eta_2}(x_2|x_1)$. By making use of the conditional independencies discussed in Section 4.2 we can often simplify these expressions further for given problems.

Using this type of approximation, the marginals $q(x_i)$ will generally be mixtures of standard distributions, which is where the added flexibility of this method comes from. By allowing the marginals $q(x_i)$ to be mixtures with dependency on the preceding elements of x , we can achieve much better approximation quality than by forcing them to be of a standard form. A practical example of this in a hierarchical Bayesian model is given in Section 8.2.

7.2 Using auxiliary variables

Another method of constructing flexible posterior approximations is by using the conditionally standard approximation of Section 7.1, but by letting the first block of variables be a vector of *auxiliary variables* z , that are not part of the unknowns x . Doing this, the posterior approximation has the form

$$q(x, z) = q(z)q(x|z).$$

The factors $q(z)$ and $q(x|z)$ should both be of standard form, which allows the marginal approximation $q(x)$ to be a general mixture of exponential family distributions, like a mixture of normals for example. If we use enough mixture components, the approximation $q(x)$ could then in principle be made arbitrarily close to $p(y|x)$. This mixture approximation can then be fitted by performing the standard KL-divergence minimization:

$$\hat{\eta} = \arg \min_{\eta} \mathbb{E}_{q_{\eta}}[\log q_{\eta}(x) - \log p(x, y)] \quad (48)$$

From (48) it becomes clear that an additional requirement of this type of approximation is that we can integrate out the auxiliary variables z from the joint $q(x, z)$ in order to evaluate the marginal density $q(x)$ at a given point x . Fortunately this is easy to do for many interesting approximations, such as discrete mixtures of normals or continuous mixtures like Student's t distributions. Also apparent from equation (48) is that we cannot use this approximation directly with the stochastic approximation algorithms proposed in the last sections since $q(x)$ is itself not part of the exponential family of distributions. However, we can rewrite (48) as

$$\hat{\eta} = \arg \min_{\eta} \mathbb{E}_{q_{\eta}}[\log q_{\eta}(x, z) - \log \tilde{p}(x, y, z)], \quad (49)$$

with $\tilde{p}(x, y, z) = p(x, y)q_{\eta}(z|x)$, and

$$q_{\eta}(z|x) = \frac{q_{\eta}(x|z)q_{\eta}(z)}{\int q_{\eta}(x|z)q_{\eta}(z)dz}.$$

Equation (49) now once again has the usual form of a KL-divergence minimization with an approximation ($q_{\eta}(x, z)$) in the exponential family. By including the auxiliary variables z in the ‘true’ posterior density, we can thus once again make use of our efficient stochastic optimization algorithms. Note that including z in the posterior did not change the marginal posterior $p(x|y)$ which is what we are interested in. A practical example of this approach, using an approximation consisting of a mixture of normals, can be found in Section 8.3.

8 Examples

We demonstrate our proposed methodology on three problems from the literature.

8.1 Binary probit regression

Binary probit (and logistic) regression is a classic model in statistics, also referred to as binary classification in the machine learning literature. Here we take a Bayesian approach to probit regression to demonstrate the performance of our methodology relative to existing variational approaches. We have N observed data pairs $(y_i \in \{0, 1\}, \mathbf{x}_i \in \mathbb{R}^P)$, and we model $y|\mathbf{x}$ as $P(y = 1|\mathbf{x}, \mathbf{w}) = \phi(\mathbf{w}'\mathbf{x})$ where $\phi(\cdot)$ is the standard Gaussian cdf and $\mathbf{w} \in \mathbb{R}^P$ is a vector of regression coefficients, for which we assume an elementwise Gaussian prior $N(0, 1)$. This is in fact a model for which existing approaches are straightforward so it is interesting to compare the performance. Of course the major benefit of our approach is that it can be applied in a much wider class of models.

We use data simulated from the model, with $N = 100$ and $P = 5$, to be able to show the performance averaged over many datasets (1000 in fact). We compare Algorithm 2 to the VBEM algorithm of Ormerod and Wand (2010) which makes use of the fact that the expectations required for this model can in fact be calculated analytically. We choose not to do this for our method to investigate how effective our MC estimation strategy can be. For completeness we also compare to variational message passing (VMP, Winn and Bishop, 2006), a message passing implementation of VBEM and expectation propagation (EP, Minka, 2001a), which is known to have excellent performance on binary classification problems (Nickisch and Rasmussen, 2008), both implemented in Infer.NET (Minka et al., 2010) a library for probabilistic inference in graphical models.

Since this experiment is on synthetic data we are able to assess performance in terms of the method’s ability to recover the known regression coefficients $\mathbf{w}$, which we quantify as the root mean squared error (RMSE) between the variational mean and the true regression weights, and the “log score”: the log density of the true weights under the approximate variational posterior. The log score is useful because it rewards a method for finding good estimates of the posterior variance as well as the mean, which should of course be central to any approximate Bayesian method.

The results, shown in Figure 2 and 3, demonstrate that our method is able to outperform the standard analytic VBEM algorithm in terms of speed accuracy tradeoff. The improvement in the RMSE is noticeable, but the difference in log score is dramatic, showing that Algorithm 2 gives significantly better estimates of the variance than VBEM. In fact our results are very similar to those of EP, which obtained an RMSE of 0.261 and log score of 0.079, but took an average of 18.2 milliseconds per run (note the system set ups are not completely comparable: EP was run on a laptop rather than a desktop, and Infer.NET is implemented in C# rather than Matlab). As expected VMP gave consistent results with VBEM: a RMSE of 0.268 and a log score of -4.85 . The average R-squared obtained by our variational approximation was 0.97, indicating a close fit to the exact posterior distribution.




Figure 2: RMSE approximate posterior mean - Stochastic Linear Regressions v.s. VBEM




Figure 3: Log-Score of approximate posterior - Stochastic Linear Regressions v.s. VBEM

8.2 A stochastic volatility model

Stochastic volatility models for signals with time varying variances are considered extremely important in finance. Here we apply our methodology to the model and prior specified in Girolami and Calderhead (2011). The data we will use, from Kim et al. (1998), is the percentage change y_t in GB Pound v.s. US Dollar exchange rate, modeled as:

$$y_t = \epsilon_t \beta \exp(v_t/2).$$

The relative volatilities, v_t are governed by the autoregressive AR(1) process

$$v_{t+1} = \phi v_t + \xi_{t+1}, \text{ with } v_1 \sim N[0, \sigma^2/(1 - \phi^2)].$$

The distributions of the error terms are given by $\epsilon_t \sim N(0, 1)$ and $\xi_t \sim N(0, \sigma^2)$. The prior specification is as in Girolami and Calderhead (2011):

$$p(\beta) \propto \beta^{-1}, \quad (\phi + 1)/2 \sim \text{Beta}(20, 1.5), \quad \sigma^2 \sim \text{Inv-Gamma}(5, 0.25)$$

Following Section 7.1 we use the hierarchical structure of the prior to suggest a hierarchical structure for the approximate posterior:

$$q_\eta(\phi, \sigma^2, \beta, v) = q_\eta(\phi)q_\eta(\sigma^2|\phi)q_\eta(\beta, v|\phi, \sigma^2).$$

The prior of ϕ is in the exponential family, so we choose the posterior approximation $q_\eta(\phi)$ to be of the same form:

$$q_\eta[(\phi + 1)/2] = \text{Beta}(\eta_1, \eta_2).$$

The prior for σ^2 is inverse-Gamma, which is also in the exponential family. We again choose the same functional form for the posterior approximation, but with a slight modification in order to capture the posterior dependency between ϕ and σ^2 :

$$q_\eta(\sigma^2|\phi) \sim \text{Inv-Gamma}(\eta_3, \eta_4 + \eta_5\phi^2),$$

where the extra term $\eta_5\phi^2$ was chosen by examining the functional form of the exact full conditional $p(\sigma^2|\phi, v)$.

The conditional prior $p[\log(\beta), v|\phi, \sigma^2]$ can be seen as the diffuse limit of a multivariate normal distribution. We therefore also use a multivariate normal conditional approximate posterior:

$$q_\eta[(\log(\beta), v)|\phi, \sigma^2] = N(m, V),$$

with

$$V^{-1} = P(\phi, \sigma^2) + \eta_6 \text{ and } m = V^{-1}\eta_7$$

where $P(\phi, \sigma^2)$ is the precision (inverse covariance) matrix of $p[(\log(\beta), v)|\phi, \sigma^2]$, η_6 is a $T \times T$ matrix, and η_7 is a $T \times 1$ vector. Furthermore, an analysis following Oppen and Archambeau (2009) shows that only a relatively small number of the elements of η_6 will be non-zero: all elements on the diagonal of η_6 and all elements in the column and row belonging to $\log(\beta)$.

Using the GB Pound vs US Dollar exchange rate data, the approximation above has almost 2000 free variational parameters to be optimized. This seems like a problematically large

number, but is easily feasible by using algorithm 2 to fit $q_\eta[\log(\beta), v|\phi, \sigma^2]$ and the algorithm using only gradients (Section 4.3) to fit $q_\eta(\phi)$ and $q_\eta(\sigma^2|\phi)$. Expectations and normalizing constants for $q_\eta[\log(\beta), v|\phi, \sigma^2]$ can be calculated efficiently using the Kalman filter and smoother (see e.g. Durbin and Koopman, 2001). For the current application we therefore only need to sample ϕ and σ^2 each iteration, and not β and v .

We compare the results against the “true” posterior, provided by a very long run of the MCMC algorithm of Girolami and Calderhead (2011).




Figure 4: Exact and approximate posterior for the stochastic volatility model - β parameter




Figure 5: Exact and approximate posterior for the stochastic volatility model - ϕ parameter




Figure 6: Exact and approximate posterior for the stochastic volatility model - σ^2 parameter

As can be seen from Figures 4, 5 and 6, the posterior approximations are nearly exact. The posterior approximation for β seems especially good (Figure 4), which is due to β being in the last block of the hierarchical posterior approximation. Similarly, the posterior approximation for the latent volatilities v (not shown) are also indistinguishable from the exact posterior.

Initializing $q_\eta(\phi)$ and $q_\eta(\sigma^2|\phi)$ to the prior, the above results can be obtained using 500 iterations of our algorithm, with a single (ϕ, σ^2) sample per iteration. Using these settings, the single-threaded Matlab implementation of our stochastic optimization algorithm requires just under a second to complete on a 3Ghz processor. This is more than two orders of magnitude faster than the running time required by advanced MCMC algorithms for this problem.

Our approach to doing inference in the stochastic volatility model shares some similarities with the approach of Liesenfeld and Richard (2008). They fit a Gaussian approximation to the posterior of the volatilities for given ϕ, σ^2, β parameters, using the importance sampling algorithm of Richard and Zhang (2007), which is based on auxiliary regressions somewhat similar to those in Algorithm 1. They then infer the model parameters using MCMC methods. The advantage of our method is that we are able to leverage the information in the gradient and Hessian of the posterior, and that our stochastic approximation algorithm allows us to fit the posterior approximation very quickly for all volatilities simultaneously, while their approach requires optimizing the approximation one volatility at a time. Unique to our approach is also the ability to concurrently fit a posterior approximation for the model parameters ϕ, σ^2, β and have the approximate posterior of the volatilities depend on these parameters, while Liesenfeld and Richard (2008) need to re-construct their approximation every time a new set of model parameters is considered. As a result, our approach is significantly faster for this problem.

8.3 A beta-binomial model for overdispersion

Albert (2009, Section 5.4) considers the problem of estimating the rates of death from stomach cancer for the largest cities in Missouri. This cancer mortality data is available from the R package LearnBayes, and consists of 20 pairs (n_j, y_j) where n_j contains the number of individuals that were at risk in city j , and y_j is the number of cancer deaths that occurred in that city. The counts y_j are overdispersed compared to what one could expect under a binomial model with constant probability, so Albert (2009) assumes the following beta-binomial model with mean m and precision K

$$P(y_j|m, K) = \binom{n_j}{y_j} \frac{B(Km + y_j, K(1 - m) + n_j - y_j)}{B(Km, K(1 - m))},$$

where $B(\cdot, \cdot)$ denotes the Beta-function. The parameters m and K are given the following improper prior

$$p(m, K) \propto \frac{1}{m(1 - m)} \frac{1}{(1 + K)^2}.$$

The resulting posterior distribution is non-standard and extremely skewed. In order to ameliorate this, Albert (2009) proposes to use the reparameterization

$$\theta_1 = \text{logit}(m), \text{ and } \theta_2 = \log(K).$$

The form of the posterior distribution $p(\theta|n, y)$ still does not resemble any standard distribution, so we will approximate it using a finite mixture of L bivariate Gaussians. In order to do this, we first introduce an auxiliary variable z , to which we assign a categorical approximate posterior distribution with L possible outcomes.

$$\log q_\eta(z) = \delta(z = 1)\eta_1 + \delta(z = 2)\eta_2 + \cdots + \delta(z = L)\eta_L,$$

where $\delta()$ is an indicator function.

Conditional on z , we then assign θ a Gaussian approximate posterior

$$q_\eta(\theta|z = i) = N(\mu_i, \Sigma_i)$$

By adapting the true posterior as described in Section 7.2, we can fit this approximate posterior to $p(\theta|n, y)$. We do this by using the basic algorithm of Section 4.1. The regression statistics C and g used in the resulting algorithm depend linearly on the indicator vector $\delta(z^* = i)$, which denotes whether or not component i was used to sample θ^* in each iteration. Rather than using this indicator function directly, we use its Rao-Blackwellized version $\mathbb{E}[\delta(z^* = i)|\theta^*]$, where θ^* are the sampled parameters. The resulting stochastic estimates will have the same expectation as when using $\delta(z^* = i)$ itself, but with lower variance at no additional computational cost.

We fit these approximations using a varying number of mixture components L and examine the resulting KL-divergence from the true posterior density. Since this is a low dimensional problem, we can obtain this divergence very precisely using quadrature methods. This also allows us to assess the accuracy of the KL-divergence approximation derived in Section 6. Finally, we present a contour plot that visually shows that a good approximation can indeed be obtained using a large enough number of mixture components.




Figure 7: KL-divergence between the variational approximation and the exact posterior density for an increasing number of mixture components.




Figure 8: Contour plots of posterior approximations using 1-8 mixture components, with the exact posterior at the bottom-right.

Figures 7 and 8 show that we can indeed approximate this skewed and fat-tailed density very well using a large enough number of Gaussians. The R-squared of the mixture approximation with 8 components is 0.997. Also apparent is the inadequacy of an approximation consisting of a single Gaussian for this problem, with an R-squared of only 0.82. This clearly illustrates the advantages of our approach which allows us to use much richer types of approximations than was previously possible. Furthermore, Figure 7 shows that the KL-divergence of the approximation to the true posterior can be approximated quite accurately using the measure developed in Section 6, especially if the posterior approximation is reasonably good.

9 Conclusion and future work

We have introduced a stochastic optimization scheme for variational inference inspired by a novel interpretation of fixed-form variational Bayes as linear regression of the target log

density against the sufficient statistics of the approximating family. Our scheme allows very generic implementation for a wide class of models since in its most basic form only the unnormalized density of the target distribution is required, although we have shown how gradient or even Hessian information can be used if necessary. The generic nature of our methodology would lend itself naturally to a software package for Bayesian inference along the lines of Infer.NET (Minka et al., 2010) or WinBUGS (Gilks et al., 1994), and would allow efficient inference in a considerably wider range of models. Incorporating automatic differentiation in such a package could clearly be beneficial. Automatic selection of the approximating family would be very appealing from a user perspective, but is currently still an open research problem.

Variational inference usually requires that we use conditionally conjugate models: since our method removes this restriction several possible avenues of research are opened. For example, collapsed versions of models (i.e. with certain parameters or latent variables integrated out) sometimes permit much more efficient MCMC inference (Porteous et al., 2008) but adapting variational methods to work with collapsed models is complex and requires custom per model methodology (Teh et al., 2007). However, our method is ambivalent to whether the model is collapsed or not, so it would be straightforward to experiment with different representations of the same model.

We have shown it is straightforward to extend our methodology to use hierarchical structured approximations and more flexible approximating families such as mixtures. This closes the gap considerably relative to MCMC methods. Perhaps the biggest selling point of MCMC methods is their asymptotic guarantees: in practice this means simply running the MCMC chain for longer can give greater accuracy, an option not available to a practitioner using variational methods. However, if we use a mixture approximating family then we can tune the computation time vs. accuracy trade off simply by varying the number of mixture components used. Another interesting direction of research along this line would be to use low rank approximating families such as factor analysis models.

It should be noted that it is possible to mix our method with VBEM, for example using our method for any non-conjugate parts of the model and VBEM for variables that happen to be conjugate. This is closely related to the non-conjugate variational message passing (NCVMP) algorithm of Knowles and Minka (2011), implemented in Infer.NET, which aims to fit conjugate models while maintaining the convenient message passing formalism. Note that NCVMP only specifies how to perform the variational optimization, not how to approximate required integrals: in Infer.NET quadrature or secondary variational bounds are used where analytic expectations are not available, unlike the Monte Carlo approach proposed here.

References

- Albert, J. (2009). *Bayesian Computation with R*. Springer Science, New York. Second edition.
- Amari, S. (1997). Neural learning in structured parameter spaces - natural Riemannian gradient. In *Advances in Neural Information Processing Systems*, pp. 127–133. MIT Press.
- Beal, M. J. and Z. Ghahramani (2002). The variational Bayesian EM algorithm for incomplete data: with application to scoring graphical model structures. In *Bayesian Statistics 7: Proceedings of the 7th Valencia International Meeting*, pp. 453.
- Bottou, L. (2010). Large-scale machine learning with stochastic gradient descent. In *Proceedings of the 19th International Conference on Computational Statistics (COMP-STAT'2010)*, pp. 177–187. Springer.
- Catcha, A. and A. Giffin (2006). Updating probabilities. In A. Mohammad-Djafari (Ed.), *Bayesian Inference and Maximum Entropy Methods in Science and Engineering*, AIP Conf. Proc.
- Durbin, J. and S. Koopman (2001). *Time Series Analysis by State Space Methods*. Oxford University Press.
- Gilks, W., A. Thomas, and D. Spiegelhalter (1994). A language and program for complex bayesian modelling. *The Statistician*, 169–177.
- Girolami, M. and B. Calderhead (2011). Riemann manifold Langevin and Hamiltonian Monte Carlo methods. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 73(2), 123–214.
- Hoffman, M., D. Blei, and F. Bach (2010). Online learning for latent Dirichlet allocation. *Advances in Neural Information Processing Systems* 23.
- Honkela, A., T. Raiko, M. Kuusela, M. Törnio, and J. Karhunen (2010). Approximate Riemannian conjugate gradient learning for fixed-form variational bayes. *Journal of Machine Learning Research*, 3235–3268.
- Kim, S., N. Shephard, and S. Chib (1998). Stochastic volatility: Likelihood inference and comparison with ARCH models. *The Review of Economic Studies* 65(3), pp. 361–393.
- Knowles, D. A. and T. P. Minka (2011). Non-conjugate variational message passing for multinomial and binary regression. In *Advances in Neural Information Processing Systems (NIPS)*.
- Liesenfeld, R. and J.-F. Richard (2008). Improving MCMC, using efficient importance sampling. *Computational Statistics and Data Analysis* 53(2), 272 – 288.

- Minka, T. P. (2001a). Expectation propagation for approximate Bayesian inference. In *Conference on Uncertainty in Artificial Intelligence (UAI)*, Volume 17.
- Minka, T. P. (2001b). *A family of algorithms for approximate Bayesian inference*. Ph. D. thesis, MIT.
- Minka, T. P., J. M. Winn, J. P. Guiver, and D. A. Knowles (2010). Infer.NET 2.4.
- Nemirovski, A., A. Juditsky, G. Lan, and A. Shapiro (2009). Robust stochastic approximation approach to stochastic programming. *SIAM J. Optim* 19(4), 1574–1609.
- Nickisch, H. and C. E. Rasmussen (2008, October). Approximations for binary Gaussian process classification. *Journal of Machine Learning Research* 9, 2035–2078.
- Opper, M. and C. Archambeau (2009). The variational Gaussian approximation revisited. *Neural Computation* 21(3), 786–792.
- Ormerod, J. T. and M. P. Wand (2010). Explaining variational approximations. *The American Statistician* 64(2), 140–153.
- Porteous, I., D. Newman, A. Ihler, A. Asuncion, P. Smyth, and M. Welling (2008). Fast collapsed Gibbs sampling for latent Dirichlet allocation. In *Proceeding of the 14th ACM SIGKDD international conference on Knowledge Discovery and Data mining*, pp. 569–577.
- Richard, J.-F. and W. Zhang (2007). Efficient high-dimensional importance sampling. *Journal of Econometrics* 141(2), 1385 – 1411.
- Robbins, H. and S. Monro (1951). A stochastic approximation method. *The Annals of Mathematical Statistics* 22(3), 400–407.
- Saul, L. and M. Jordan (1996). Exploiting tractable substructures in intractable networks. *Advances in Neural Information Processing Systems*, 486–492.
- Storkey, A. J. (2000). Dynamic trees: A structured variational method giving efficient propagation rules. In *Conference on Uncertainty in Artificial Intelligence (UAI)*.
- Teh, Y., D. Newman, and M. Welling (2007). A collapsed variational Bayesian inference algorithm for latent Dirichlet allocation. *Advances in Neural Information Processing Systems* 19, 1353.
- Turner, R. E., P. Berkes, and M. Sahani (2008). Two problems with variational expectation maximisation for time-series models. *Inference and Estimation in Probabilistic Time-Series Models*.
- Winn, J. and C. M. Bishop (2006). Variational message passing. *Journal of Machine Learning Research* 6(1), 661.

A Bias and variance calculations

We first consider the bias of the estimators. Consider the multivariate Taylor expansion of $f : \mathbb{R}^K \rightarrow \mathbb{R}$ around the point $\bar{y} \in \mathbb{R}^K$:

$$f(y) \approx f(\bar{y}) + (y - \bar{y})' f'(\bar{y}) + \frac{1}{2} \text{tr}((y - \bar{y})(y - \bar{y})' \nabla^2 f(\bar{y})) \quad (50)$$

From this we can derive expressions for the expectation of $f(y)$:

$$\mathbb{E}f \approx f(\bar{y}) + \frac{1}{2} \text{tr}(\text{Cov}(y) f''(\bar{y})) \quad (51)$$

where we have chosen to perform the Taylor expansion around the mean $\bar{y} = \mathbb{E}y$. For the first estimator let $y = \frac{1}{S} \sum_s a(x_s)$ and $f(y) = 1/y$, then we find

$$\mathbb{E}\hat{\eta}_1 = \mathbb{E} \left[\left(\frac{1}{S} \sum_s a(x_s) \right)^{-1} \right] \mathbb{E}[b] \quad (52)$$

$$\approx \left(\frac{1}{\mathbb{E}[a]} + \frac{\text{Var}(a)}{S \mathbb{E}[a]^3} \right) \mathbb{E}[b] \quad (53)$$

$$= \mathbb{E}[\eta] + \frac{\text{Var}(a) \mathbb{E}[b]}{S \mathbb{E}[a]^3} \quad (54)$$

since $\text{Var}(y) = \text{Var}(a)/S$. We see that the bias term depends on the ratio $\text{Var}(a)/\mathbb{E}[a]^2$, i.e. the spread of the distribution of a relative to its magnitude.

Now for the second estimator let

$$y = \begin{bmatrix} \frac{1}{S} \sum_s a(x_s) \\ \frac{1}{S} \sum_s b(x_s) \end{bmatrix} \quad (55)$$

so that $\eta_2 = f(y) = \frac{y_2}{y_1}$. Note that $\text{Cov}(y) = \frac{1}{S} \text{Cov}([a, b]')$ and

$$\nabla^2 f(y) = \begin{bmatrix} \frac{2y_2}{y_1^3} & -\frac{1}{y_1^2} \\ -\frac{1}{y_1^2} & 0 \end{bmatrix} \quad (56)$$

Putting everything together we have

$$\mathbb{E}\hat{\eta}_2 \approx \mathbb{E}\eta + \frac{\text{Var}(a) \mathbb{E}b}{S \mathbb{E}[a]^3} - \frac{\text{Cov}(a, b)}{S \mathbb{E}[a]^2} \quad (57)$$

Note that we recover the expression for $\mathbb{E}\hat{\eta}_1$ if $\text{Cov}(a, b) = 0$, which makes sense because if we use different randomness for calculating $\mathbb{E}[a]$ and $\mathbb{E}[b]$ then a, b have 0 covariance in our MC estimate. Finally the analytic estimator is unbiased:

$$\mathbb{E}\hat{\eta}_a = \mathbb{E}\eta \quad (58)$$

We now turn to the variances. The analytic estimator is a standard MC estimator with variance

$$\text{Var}(\hat{\eta}_a) = \frac{\text{Var}(b)}{S\mathbb{E}(a)^2} \quad (59)$$

Consider only the linear terms of the Taylor expansion:

$$f(y) \approx f(\bar{y}) + (y - \bar{y})' f'(\bar{y}) \quad (60)$$

Substituting this into the formula for variance gives

$$\text{Var}[f(y)] = \mathbb{E}[(f(y) - \mathbb{E}f(y))(f(y) - \mathbb{E}f(y))'] \quad (61)$$

$$\approx \mathbb{E}[f'(\bar{y})'(y - \bar{y})(y - \bar{y})' f'(\bar{y})] \quad (62)$$

$$= f'(\bar{y})' \text{Var}(y) f'(\bar{y}) \quad (63)$$

We will calculate the variance of the second estimator and derive the variance of the first estimator from this. Again let y be as in (55). Note that $\text{Var}(y) = \text{Cov}(a, b)/S$. We find

$$\text{Var} \hat{\eta}_2 \approx \frac{1}{S} \left(\frac{(\mathbb{E}b)^2 \text{Var} a}{(\mathbb{E}a)^4} - 2 \frac{\mathbb{E}b \text{Cov}(a, b)}{(\mathbb{E}a)^3} + \frac{\text{Var} b}{(\mathbb{E}a)^2} \right) \quad (64)$$

The final term is equal to that for the analytic estimator. The second term is not present in the variance of the first estimator, since then a and b have no covariance under the sampling distribution, i.e.

$$\text{Var} \hat{\eta}_1 \approx \frac{1}{S} \left(\frac{(\mathbb{E}b)^2 \text{Var} a}{(\mathbb{E}a)^4} + \frac{\text{Var} b}{(\mathbb{E}a)^2} \right) \quad (65)$$

The first term is always positive, suggesting that $\hat{\eta}_1$ is dominated by the analytic estimator. If we make the assumption that $p(x)$ is in the same family as q then we have $\log p(x) = \lambda T(x)$, then we find

$$\mathbb{E}a = \mathbb{E}[T^2] \quad (66)$$

$$\mathbb{E}b \approx \lambda \mathbb{E}[T^2] \quad (67)$$

$$\text{Cov}(a, b) \approx \lambda \mathbb{E}[T^4] - \lambda \mathbb{E}[T^2]^2 = \eta \text{Var}(T^2) \quad (68)$$

$$\text{Var}(a) = \mathbb{E}[T^4] - \mathbb{E}[T^2]^2 = \text{Var}(T^2) \quad (69)$$

B Unnormalized to normalized optimality condition

The unnormalized optimality condition in (7) is

$$\tilde{\eta} = \left[\int \tilde{q}_{\tilde{\eta}}(x) \tilde{T}(x)' \tilde{T}(x) d\nu(x) \right]^{-1} \left[\int \tilde{q}_{\tilde{\eta}}(x) \tilde{T}(x)' \log p(x, y) d\nu(x) \right]. \quad (70)$$

Clearly we can replace $\tilde{q}$ by its normalized version $q(x) = \tilde{q}/Z(\eta)$ since the normalizing terms will cancel. Recalling $\tilde{T}(x) = (1, T(x))$ and $\tilde{\eta} = (\eta_0, \eta)'$ we then have

$$\begin{bmatrix} 1 & \mathbb{E}T \\ \mathbb{E}T' & \mathbb{E}[T'T] \end{bmatrix}^{-1} \begin{pmatrix} \mathbb{E}Y \\ \mathbb{E}[TY] \end{pmatrix} = \begin{pmatrix} \eta_0 \\ \eta \end{pmatrix} \quad (71)$$

where $Y := \log p(x, y)$. Rearranging gives

$$\begin{pmatrix} \mathbb{E}Y \\ \mathbb{E}[TY] \end{pmatrix} = \begin{bmatrix} 1 & \mathbb{E}T \\ \mathbb{E}T' & \mathbb{E}[T'T] \end{bmatrix} \begin{pmatrix} \eta_0 \\ \eta \end{pmatrix} \quad (72)$$

Solving for η_0 easily gives

$$\eta_0 = \mathbb{E}Y - \mathbb{E}[T]\eta \quad (73)$$

$$\eta = (\mathbb{E}[T'T] - \mathbb{E}T'\mathbb{E}T)^{-1} (\mathbb{E}[TY] - \mathbb{E}T'\mathbb{E}Y) \quad (74)$$

$$= \text{Cov}(T)^{-1} \text{Cov}(T, Y) \quad (75)$$

C Derivation of Gaussian variational approximation

For notational simplicity we will derive our stochastic approximation algorithm for Gaussian variational approximation (algorithm 2) under the assumption that x is univariate. The extension to multivariate x is conceptually straightforward but much more tedious in terms of notation.

Let $p(x, y)$ be the unnormalized posterior distribution of a univariate random variable x , and let $q(x) = N(m, V)$ be its Gaussian approximation. In order to find the mean m and variance V that minimize the KL-divergence between $q(x)$ and $p(x|y)$ we solve the transformed regression problem defined by

$$K(\eta) = [\nabla_{\eta}\phi(\eta)]^{-1},$$

with $\phi = (m, V)'$ the usual mean-variance parameterization and where the natural parameters are given by $\eta = (V^{-1}m, V^{-1})'$. Following equation (43), we know that the regression statistics for this optimization problem are given by

$$K(\eta) \text{Cov}_{q_{\phi}}[T(x), T(x)] = \nabla_{\phi}\mathbb{E}_{q_{\phi}}T(x) \quad (76)$$

with $T(x) = (x, -0.5x^2)'$ the vector of sufficient statistics, and

$$K(\eta) \text{Cov}_{q_{\phi}}[T(x), \log p(x, y)] = \nabla_{\phi}\mathbb{E}_{q_{\phi}}\log p(x, y). \quad (77)$$

Recall identity (39) which states that

$$\nabla_{\phi_1} \mathbb{E}_{q_\phi} g(x) = \mathbb{E}_{q_\phi} \nabla_x g(x),$$

with $\phi_1 = m$ the first element of the parameter vector ϕ , and $g(x)$ any differentiable function. Similarly, identity (40) reads

$$\nabla_{\phi_2} \mathbb{E}_{q_\phi} g(x) = -\frac{1}{2} \mathbb{E}_{q_\phi} \nabla_x \nabla_x g(x),$$

with $\phi_2 = V$ the second element of the parameter vector. Using these identities to stochastically approximate the regression statistic (77) we obtain

$$\hat{g}_1 = \nabla_x \log p(x, y)|_{x=x^*}, \text{ and } \hat{g}_2 = -\frac{1}{2} \nabla_x \nabla_x \log p(x, y)|_{x=x^*},$$

with x^* a draw from $q_\phi(x)$. We can approximate the regression statistic (76) in a similar way. Making use of the fact that $T(x) = (x, -0.5x^2)'$ is the vector of sufficient statistics, we obtain

$$\begin{aligned} \hat{C}_{1,1} &= 1 \\ \hat{C}_{1,2} &= -x^* \\ \hat{C}_{2,1} &= 0 \\ \hat{C}_{2,2} &= -0.5. \end{aligned} \tag{78}$$

Note that $\hat{C}_{1,2}$ is the only stochastic entry in the matrix, which means we only have to store and average the samples x^* , rather than the full C matrix. We will denote the average of these samples by $\bar{z}$. Storing the 2-dimensional $\hat{g}$ statistic is equivalent to storing the averaged gradient and Hessian of $\log p(x, y)$, which we denote by $\bar{a}$ and $\bar{P}$. Since our estimate $\hat{C}$ is upper-diagonal, the final regression step $\bar{\eta} = \bar{C}^{-1} \bar{g}$ from algorithm 1 can easily be expressed analytically in terms of these quantities, which gives $V = \bar{P}^{-1}$ and $m = V \bar{a} + \bar{z}$ as the optimal mean and variance of the approximation.

D Stochastic linear regression and gradient descent

At each iteration of Algorithm 1, the current variational parameters η_t are stochastically updated to a new value η_{t+1} . In Section 3 we derive this update from the viewpoint of linear regression. In this section we show that, for very small step-sizes, the update is also similar to a preconditioned stochastic gradient descent step. As in Algorithm 1, we have $\eta_t = C_t^{-1} g_t$, and we update this to

$$\eta_{t+1} = [(1 - w_t)C_t + w_t\hat{C}_t]^{-1}[(1 - w_t)g_t + w_t\hat{g}_t] = [C_t + \lambda_t\hat{C}_t]^{-1}[g_t + \lambda_t\hat{g}_t],$$

where $\hat{g}_t$ and $\hat{C}_t$ are the stochastic estimates generated during iteration t , and where $\lambda_t = w_t/(1 - w_t)$ is the effective step-size of the update. To characterize this update for small values of λ_t we perform a first order Taylor expansion of η_{t+1} around $\lambda_t = 0$, which gives

$$\eta_{t+1} = \eta_t - \lambda_t C_t^{-1}(\hat{C}_t \eta_t - \hat{g}_t) + \mathcal{O}(\lambda_t^2). \quad (79)$$

Comparison with equation (6) shows that the stochastic term in this expression ($\hat{C}_t \eta_t - \hat{g}_t$) is an unbiased estimate of the gradient of the KL-divergence $D[q_{\eta_t}(x)|p(x, y)]$. Up to first order, the update equation in (79) thus represents a stochastic gradient descent step, preconditioned with the C_t^{-1} matrix. Since this pre-conditioner is independent of the stochastic gradient approximation at iteration t , this gives a valid adaptive stochastic gradient descent algorithm, to which all the usual convergence results apply (see e.g. Amari, 1997). Note the connection between this result and the result of equation (18): there we show that the bias disappears as the number of samples grows large, while here we show that it disappears as the step-size gets small. In Section 3 we further show that the effect of this bias on the update η_{t+1} disappears faster than the effect of the variance.

If we take small steps, the pre-conditioner C_t^{-1} of equation (79) will be close to the Riemannian metric $\mathbb{E}_{q_t} \hat{C}_t = \text{Var}_{q_t}[T(x)]$ used in natural gradient descent algorithms like that of Honkela et al. (2010). For certain exponential family distributions this metric can be calculated analytically, which suggests performing natural gradient descent optimization with updates of the form

$$\eta_{t+1} = \eta_t - \lambda_t \left(\eta_t - \text{Var}_{q_t}[T(x)]^{-1} \text{Cov}_{q_t}[T(x), \log p(x, y)] \right),$$

where the covariance $\text{Cov}_{q_t}[T(x), \log p(x, y)]$ is approximated using Monte Carlo. While this approach is clearly related to Algorithm 1 there are two significant differences: 1) this would correspond closely to the analytic estimator in (12), which we have seen can both theoretically and empirically be improved on by using the linear regression estimator (11). This means that the $\mathcal{O}(\lambda_t^2)$ term in equation (79) is generally beneficial. 2) Algorithm 1 efficiently averages the MC samples over multiple iterations so that just a single MC sample per iteration is required. This is statistically more efficient than performing averaging in the space of the η parameters, as is done in stochastic gradient descent.